

SHOCK®

Injury, Inflammation, and Sepsis: Laboratory and Clinical Approaches

OFFICIAL JOURNAL OF THE SHOCK SOCIETY, THE EUROPEAN SHOCK SOCIETY, THE INDONESIAN SHOCK SOCIETY, THE INTERNATIONAL FEDERATION OF SHOCK SOCIETIES, AND THE OFFICIAL AND INTERNATIONAL JOURNAL OF THE JAPAN SHOCK SOCIETY

Volume 33, No. 5

May 2010

<i>Kevin J. Tracey</i>	451	<u>Commentary</u> What's New in <i>Shock</i>, May 2010?
<hr/>		
<i>Chih-Hsin Lee, Jann-Yuan Wang, Yao-Kuang Wu, Hung-Wen Chiu, Chou-Chin Lan, Hung Chang, and Chi-Yuan Chen</i>	454	<u>Clinical and Basic Science Aspects</u> Pulse Pressure Power Spectrum Predicts Volume Responsiveness in Shock Patients Without Sedation Supplemental Digital Content is available in the text.
<hr/>		
<i>Anubhav Mittal, Friederike Göke, Richard Flint, Benjamin P.T. Loveday, Nichola Thompson, Brett Delahunt, Paul A. Kilmartin, Garth J.S. Cooper, Julia MacDonald, Anthony Hickey, John A. Windsor, and Anthony R.J. Phillips</i>	460	<u>Basic Science Aspects</u> The Redox Status of Experimental Hemorrhagic Shock as Measured by Cyclic Voltammetry
<i>Ahmad W. Husari, Aline Khayat, Haitham Awdeh, Hadi Hatoum, Michel Nasser, Salman M. Mroueh, Ghazi Zaatari, Marwan El-Sabban, and Ghassan S. Dbaibo</i>	467	Activated Protein C Attenuates Acute Lung Injury and Apoptosis in a Hyperoxic Animal Model
<i>Shiu-Jen Chen, Shaio-Yun Li, Chih-Chin Shih, Mei-Huei Liao, and Chin-Chen Wu</i>	473	NO Contributes to Abnormal Vascular Calcium Regulation and Reactivity Induced by Peritonitis-Associated Septic Shock in Rats
<i>Elif Cadirci, Berrin Zuhul Altunkaynak, Zekai Halici, Fehmi Odabasoglu, M. Hamidullah Uyanik, Cemal Gundogdu, Halis Suleyman, Mesut Halici, Mevlut Albayrak, and Bunyami Unal</i>	479	α-Lipoic Acid as a Potential Target for the Treatment of Lung Injury Caused by Cecal Ligation and Puncture-Induced Sepsis Model in Rats
<i>Huai-Chia Chang, Ke-Hsun Lin, Yu-Ting Tai, Juei-Tai Chen, and Ruei-Ming Chen</i>	485	Lipoteichoic Acid-Induced <i>TNF-α</i> and <i>IL-6</i> Gene Expressions and Oxidative Stress Production in Macrophages are Suppressed by Ketamine Through Downregulating Toll-Like Receptor 2-Mediated Activation of ERK1/2 and NFκB
<i>Nandini Verma, Subhash K. Tripathi, Indrajit Chaudhury, Hasi R. Das, and Rakha H. Das</i>	493	iNOS-Targeted 10-23 DNzyme Reduces LPS-Induced Systemic Inflammation and Mortality in Mice
<i>Nobuaki Kawarabayashi, Shuhji Seki, Kazuo Hatsuse, Manabu Kinoshita, Toshimichi Takigawa, Hironori Tsujimoto, Toshinobu Kawabata, Hiroyuki Nakashima, Satoshi Shono, and Hidetaka Mochizuki</i>	501	Immunosuppression in the Livers of Mice with Obstructive Jaundice Participates in Their Susceptibility to Bacterial Infection and Tumor Metastasis
<i>Jan P. Roesner, Jan Mersmann, Stefan Bergt, Karl Bohnenberg, Carmen Barthuber, Csaba Szabo, Gabriele E. F. Nöldge-Schomburg, and Kai Zacharowski</i>	508	Therapeutic Injection of PARP Inhibitor INO-1001 Preserves Cardiac Function in Porcine Myocardial Ischemia and Reperfusion Without Reducing Infarct Size

<i>Uta Kerkweg, Katja Pamp, Judith Fieker, Frank Petrat, Robert C. Hider, and Herbert de Groot</i>	514	Release of Redox-Active Iron by Muscle Crush Trauma: No Liberation Into the Circulation
<i>Isabelle Chapados, Constance L. Chik, and Po-Yin Cheung</i>	520	Plasma Cortisol Response to ACTH Challenge in Hypoxic Newborn Piglets Resuscitated with 21% and 100% Oxygen
<i>Koichi Suda, Hiroya Takeuchi, Tomoko Hagiwara, Taku Miyasho, Minoru Okamoto, Kazufumi Kawasaki, Shingo Yamada, Kazuhiro Saganuma, Norihito Wada, Yoshiro Saikawa, Koichi Fukunaga, Yosuke Funakoshi, Satoru Hashimoto, Hiroshi Yokota, Ikuro Maruyama, Akitoshi Ishizaka, and Yuko Kitagawa</i>	527	Neutrophil Elastase Inhibitor Improves Survival of Rats with Clinically Relevant Sepsis
<i>Pushpa Sharma and Paul D. Mongan</i>	533	Hypertonic Sodium Pyruvate Solution is More Effective Than Ringer's Ethyl Pyruvate in the Treatment of Hemorrhagic Shock
<i>Rosanna Di Paola, Alessia Melani, Emanuela Esposito, Emanuela Mazzon, Irene Paterniti, Placido Bramanti, Felicita Pedata, and Salvatore Cuzzocrea</i>	542	Adenosine A_{2A} Receptor-Selective Stimulation Reduces Signaling Pathways Involved in the Development of Intestine Ischemia and Reperfusion Injury
<i>Editorial Comment</i>		
<i>Sebastian Rehberg, Christoph Thiemermann, and Marc O. Maybauer</i>	553	Role of Chelatable Iron Versus Myoglobin in Oxidative Stress After Crush Trauma
<i>Letters to the Editor</i>		
<i>To the Editor: Lutz Schomburg</i>	555	A Large-Bolus Injection, But Not Continuous Infusion of Sodium Selenite Improves Outcome in Peritonitis.
<i>Reply: Zhen Wang, Jean-Louis Vincent, and Xavier Forceville</i>		
<i>To the Editor: Pavlos M. Myrianthefs, Eleni Boutzouka, and George J. Baltopoulos</i>	557	γ-Globulin Levels in Patients with Community-Acquired Septic Shock
<i>Reply: Fabio Silvio Taccone, Daniel De Backer, and Jean-Louis Vincent</i>		

SHOCK® is abstracted and/or indexed in *Index Medicus*, MEDLINE, Current Contents®/Life Sciences, Science Citation Index®, SciSearch®, Research Alert®, the Biochemistry & Biophysics Citation Index™, and Reference Update Current Impact Factor 3.394

Instructions for Authors are available online at http://journals.lww.com/shockjournal/Documents/SHOCK_IFA.pdf and are printed in the June and December issues of the journal.

COVER: Immunohistochemical staining of NF- κ B (p65) in lung tissue sections of CLP-operated and α -lipoic acid (ALA)-administered rats. The minimal positivity was especially seen in the cytoplasmic pattern of only the interalveolar septa. Sham control (C, D); ALA-administered rats (A, B). See Cadirci et al., pages 479–484, 2010.